ТАКРОРИЙ ЭКИН СИФАТИДА ЭКИЛГАН СОЯ НАВЛАРИНИНГ МАЙСА ДАВРИДАГИ МОРФОЛОГИК КЎРСАТКИЧЛАРИ

М. А. Холиқова
ТВЧДПИ
holiqova19811@mail.ru

Ҳ. Х. Матниязова
Генетика ва ўсимликлар экспериментал биологияси институти
matniyazova@mail.ru

Г. С. Исмагилова
ТВЧДПИ
gulnozaismagilova@gmail.com

АННОТАЦИЯ

Майсалаш даврида соянинг илдиз тизими ривожланади. Ушбу ўсимлик тўлиқ майсалаш даврга кирганда унинг илдизида туганаклар ривожланади. Натижада, ризобиум бактериялари ҳаво азотини ўзлаштириб, ўсимликни азот билан таъминлайди ва бир қисм азотни тупроқда қолдириб, унинг унумдорлигини оширади.

Калит сўзлар: соя, такрорий экин, морфохўжалик, майса, илдиз.

MORPHOLOGICAL INDICATIONS OF SOY VARIETIES PLANTED AS REPRODUCTIVE CROPS

M. A. Kholikova
Teacher of CSPI
holiqova19811@mail.ru

H. X. Matniyazova
Institute of Genetics and Experimental Biology of Plants
textiyazova@mail.ru

G. S. Ismagilova
Teacher of CSPI
gulnozaismagilova@gmail.com

ABSTRACT

During mowing, the root system of shade develops. When this plant enters the period of full weeding, it develops buds on its roots. As a result, rhizobium bacteria assimilate air nitrogen, providing the plant with nitrogen and leaving some nitrogen in the soil, increasing its fertility.

Keywords: soy, replanting, morphology, grass, root