АСОСИЙ ЭКИН СИФАТИДА ЭКИЛГАН АЙРИМ МАҲАЛЛИЙ ВА ХОРИЖИЙ СОЯ НАВЛАРИНИНГ ҲОСИЛДОРЛИК КЎРСАТКИЧЛАРИ

М. А. Холиқова
ТВЧДПИ
holiqova19811@mail.ru

Ҳ. Х. Матниязова
Генетика ва ўсимликлар экспериментал биологияси институти, ТВЧДПИ
matniyazova@mail.ru

АННОТАЦИЯ

Ушбу мақолада Самарқанд вилоятида асосий экин сифатида экилган соя навларининг муҳим морфохўжалик белгиларидан бири бўлган битта ўсимликдаги ҳосилдорлик кўрсаткичи ўрганилди. Олинган маълумотларга кўра хорижий навларда Қозоғистон навларидан Нена нави, Россия навларидан Спарта нави ва маҳаллий навлардан Орзу навида дуккаклаш ва 1000 та дон оғирлиги бўйича юқори кўрсаткич қайд этилди.

Калит сўзлар: соя, асосий экин, хўжалик, дуккак

FERTILITY INDICATORS OF SOME LOCAL AND FOREIGN SOY VARIETIES PLANTED AS MAIN CROPS

M. Kholiqova
Teacher of CSPI

H. X. Matniyazova
Institute of Genetics and Experimental Biology of Plants

ABSTRACT

In this article, the yield index of a single plant, which is one of the important morphological features of soybean varieties planted as the main crop in Samarkand region, was studied. According to the data, foreign varieties have a high yield of Nena from Kazakhstan, Sparta from Russia and Orzu from local varieties.

Keywords: shade, main crop, farm, legume.