БАЧАДОН БЎЙНИ ХАВФЛИ ЎСМАЛАРИНИНГ ЁШГА ХОС ХУСУСИЯТЛАРИ

Г. Ф. Махмудова
Бухоро давлат тиббиёт институти

Д. В. Темирова
Бухоро давлат тиббиёт институти

Ш. Б. Баротова
Бухоро давлат тиббиёт институти

АННОТАЦИЯ

Тадқиқотнинг мақсади фертил ва кекса ёшдаги аёлларда бачадон бўйни хавфли ўсмаларининг клинико-анатомо-морфологик хусусиятларини ўрганишдан иборат. Изланиш бачадон бўйни хавфли ўсмаси бўлган 60 нафар беморнинг ретроспектив ва проспектив маьлумотлари асосида олиб борилди. Касалланиш 52,2 ёшли аёлларда 35-39 ёш хамда 60-64 ёшлиларга нисбатан кўпрок кузатилмокда. Айтиш мумкинки, аёлларда 83,3% холатларда ясси хужайрали рак тури учрамокда. Айнан катта ёшли аёлларда ракнинг бу тури эндофит шаклда ўсиб давонинг қийинлашига ва прогнозининг яхши бўлмаслигига сабаб бўлмокда.

Калит сўзлар: эндофит, ретроспектив, проспектив, ясси хужайрали рак.

AGE-RELATED SPECIFIC FEATURES OF MALIGNANT TUMORS OF THE CERVIX

ABSTRACT

The aim of the study is to study the clinical, anatomical and morphological features of malignant neoplasms of the cervix uteri in fertile and elderly women. The study is based on retrospective and prospective data from 60 patients with malignant cervical cancer. The disease is more common in 52.2-year-old women than in 35-39-year-olds and 60-64-year-olds. We can say that in 83.3% of cases, women develop squamous cell carcinoma. It is in older women that this type of cancer develops into an endophyte, which complicates treatment and worsens the prognosis.

Keywords: endophyte, retrospective, prospect, squamous cell carcinoma.