# ДИСК ЧУРРАЛАРИ РИВОЖЛАНГАН БЎЙИН ОСТЕОХОНДРОЗЛАРИНИНГ ЭТИОПАТОГЕНЕЗИ ВА КЛИНИК МАНЗАРАСИ

**Зухриддин Амриддин ўғли Примов**
Самарқанд давлат тиббиёт институти магистратура резиденти

**Даврон Мавлонович Равшанов**
Самарқанд давлат тиббиёт институти мустақил изланувчиси

**Амирқул Шодиевич Шодиев**
Самарқанд давлат тиббиёт институти доценти, тиббиёт фанлари доктори

## АННОТАЦИЯ

Мақолада бўйин остеохондрозида ривожланадиган умуртқалараро диск чурраси билан даволанган 62 та беморда касаллик этиопатогенези ва клиник-неврологик манзараси ўрганилган. Диск чурраларининг келиб чиқишида жисмоний зўриқиш(46,0%), кам ҳаракатлик ва ортиқча вазн(35,4%), травмалар (4,8%) сабаб бўлади. Бўйин соҳаси диск чурралари қўлларда сезгининг бузилиши (93,5%), елка ва қўллардаги оғриқ(90,3%), мушаклар гипотрофияси (87,1%), бош айланиши(72,6%), қўллардаги уюшишлар(69,3%), бош оғриғи (66,1%), сингари клиник симптомлар билан намоён бўлади.

**Калит сўзлар:** Диск чурраси, умуртқа, гипотрофия, облитерация, синдром.

# FEATURES OF ETIOPATHOGENESIS AND CLINICAL COURSE OF OSTEOCHONDROSIS OF THE SPINE WITH HERNIATED INTERVERTEBRAL DISCS

## ABSTRACT

In this article studied the etiopathogenesis and clinical manifestation of intervertebral herniated discs are sing in cervical osteochondrosis of the spine in 62 patients. Physical overstrain(46,0%), low mobility and excess weight(35,4%), trauma (4,8%) are the main causes of disc herniations. Intervertebral disc herniations in the cervical spine are manifested by clinical symptoms such as decrease of superficial sensation in hands(93,5%), pain in the shoulder and hands(90,3%), muscle hypotrophy(87,1%), dizziness(72,6%), numbness in hands(69,3%), headache(66,1%), and etc.

**Keywords:** disc herniation, hypotrophy, obliteration.